

**Negative scattering asymmetry parameter for dipolar particles:  
Unusual reduction of the transport mean free path and radiation pressure**

R. Gómez-Medina,<sup>1</sup> L. S. Froufe-Pérez,<sup>1</sup> M. Yépez,<sup>1</sup> F. Scheffold,<sup>2</sup>  
M. Nieto-Vesperinas,<sup>3</sup> and J. J. Sáenz,<sup>1</sup>

<sup>1</sup> Dpto. Física de la Materia Condensada and Instituto Nicolás Cabrera,

Universidad Autónoma de Madrid, Campus de Cantoblanco, Madrid 28049, Spain.

<sup>2</sup> Department of Physics, University of Fribourg, Chemin du Musé 3, 1700 Fribourg, Switzerland

<sup>3</sup> Instituto de Ciencia de Materiales de Madrid, Consejo Superior de Investigaciones Científicas (CSIC),  
Campus de Cantoblanco, Madrid 28049, Spain.  
[r.gomezmedina@uam.es](mailto:r.gomezmedina@uam.es)

Propagation of light and image formation in turbid media has long been a subject of great interest [1] and constitutes the core of powerful techniques with countless applications including biomedical imaging [2, 3] and dynamic spectroscopy techniques [4-6], characterization of composite materials and complex fluids [7], remote sensing or telecommunications [8] to mention a few.

Lossless dielectric nanospheres (made of nonmagnetic materials) with relatively low refraction index may present strong electric and magnetic dipolar resonances [9-11].

We establish a relationship between the optical force [12,13] from a plane wave on small electric and magnetic dipolar particles, the transport cross section, and the scattering asymmetry parameter  $g$  [14].


In this way we predict negative  $g$  (that minimize the transport mean free path below values of the scattering mean free path) for a dilute suspension of both perfectly reflecting spheres as well as of lossless dielectric nanospheres made of moderate permittivity materials, e.g., silicon or germanium nanospheres in the infrared region. Lossless dielectric Mie spheres of relatively low refraction index (as low as 2.2) are shown to present negative  $g$  in specific spectral ranges [14].

## References

- [1] *Waves and Imaging through Complex Media*, edited by P. Sebbah (Kluwer Academic, Dordrecht, 2001); in *Wave Scattering in Complex Media: From Theory to Applications*, edited by B. A. van Tiggelen and S. E. Skipetrov, NATO Science Series II: Mathematics, Physics and Chemistry, Vol. 107 (Kluwer Academic, Dordrecht, 2003).
- [2] A. Yodh and B. Chance, Phys. Today **48**(3), 34 (1995); S. K. Gayen and R. R. Alfano, Opt. Photon. News **7** (1996) 17.
- [3] J. Ripoll, V. Ntziachristos, J. P. Culver, D. N. Pattanayak, A. G. Yodh, and M. Nieto-Vesperinas, J. Opt. Soc. Am. A **18** (2001) 821.
- [4] D. A. Weitz and D. J. Pine, in *Dynamic Light Scattering*, edited by W. Brown (Oxford University Press, New York, 1993).
- [5] G. Maret and P. E. Wolf, Z. Phys. B **65** (1987) 409; D. J. Pine, D. A. Weitz, P. M. Chaikin, and E. Herbolzheimer, Phys. Rev. Lett. **60** (1988) 1134.
- [6] R. Lenke and G. Maret, in *Multiple Scattering of Light: Coherent Backscattering and Transmission*, edited by W. Brown (Gordon & Breach, Reading, UK, 2000).
- [7] F. Scheffold and P. Schurtenberger, Soft Mater. **1** (2003) 139.
- [8] A. Derode, A. Tourin, J. de Rosny, M. Tanter, S. Yon, and M. Fink, Phys. Rev. Lett. **90** (2003) 14301; G. Lerosey, J. de Rosny, A. Tourin, and M. Fink, Science **315** (2007) 1120.

- [9] A. García-Etxarri, R. Gómez-Medina, L. S. Froufe-Pérez, C. López, L. Chantada, F. Scheffold, J. Aizpurua, M. Nieto-Vesperinas and J. J. Sáenz, *Opt. Express*, **19** (2011) 4815-4826.
- [10] R. Gómez-Medina, B. García-Cámarra, I. Suárez-Lacalle, F. González, F. Moreno, M. Nieto-Vesperinas and J. J. Sáenz, *J. Nanophoton.* **5** (2011) 053512.
- [11] M. Nieto-Vesperinas, R. Gómez-Medina, and J. J. Sáenz, *J. Opt. Soc. Am. A*, **28** (2011) 54-60.
- [12] M. Nieto-Vesperinas, J. J. Sáenz, R. Gómez-Medina, and L. Chantada, *Opt. Express*, **18** (2010) 11428-11443.
- [13] R. Gómez-Medina, M. Nieto-Vesperinas and J. J. Sáenz, *Phys. Rev. A*, **83** (2011) 033825.
- [14] R. Gómez-Medina, L. S. Froufe- Pérez, M. Yépez, F. Scheffold, M. Nieto-Vesperinas and J. J. Sáenz, *Phys. Rev. A*, **85** (2012) 035802.

## Figures



**Figure 1.** (a) Color map of the  $g$  factor for spherical absorptionless particles as a function of their refractive index  $m$  and size parameter  $y = mka$ . As seen in the attached scale, green areas correspond to negative values of  $g$ . (b) Color map of the sphere scattering cross section. Red corresponds to dominant electric dipole contributions to the scattering cross section. Green corresponds to dominant magnetic dipole contributions, while blue sums up all higher-order multipole terms. Vertical dashed lines coincide with  $y$  parameter for maximum electric dipole contribution (right vertical line) and maximum magnetic dipole contribution (left vertical line). The white horizontal line at  $m \approx 3.5$  which corresponds to a silicon sphere. (After Ref.[14]).